

Key 1 – Key to Insects Orders


Notes:

- This key covers insect orders commonly and occasionally observed. However, it does not include all orders. Key #1 is similar, but easier, being limited to insect orders commonly associated with gardening and landscape maintenance.
 - This simplified key covers insects showing common characteristics for the order. Species with atypical appearances will not work with this simplified key.
-

1a. Wings present. — go to 2

1b. Wings absent. — go to 9

(1a) 2a. One pair of membranous wings. The insect may look like a fly or bee. (Look carefully for a second pair of wings hidden beneath the front pair.) — *Diptera*(flies)



2b. Two pair of wings present. — go to 3

(2b) 3a. Front and hind wings not similar in texture. Front wings parchment-like, shell-like or leathery or thickened. Hind wings more delicate or membranous. — go to 4

3b. Front and hind wings similar in texture. Either membranous, transparent or covered with powdery-like scales. — go to 7


(3a) 4a. Chewing mouthparts. — go to 5



4b. Piercing-sucking mouthparts. — go to 6




(4a) 5a. Leathery front wings, without veins, lay over body in a shell-like fashion, making a straight line between wings. Hind wings membranous. No cerci (tail-like appendages). — **Coleoptera** (beetle)




5b. Front wings more or less parchment like with a network of veins. Hinds wings membranous, usually broad with folds like a fan. Hind leg enlarged for jumping — **Orthoptera** (grasshoppers, crickets, katydids)




5c. Front wings more or less parchment like with a network of veins. Flattened bodies, thread-like antennae — **Blattaria** (cockroach)




5d. Front wings more or less parchment like with a network of veins. Front legs modified to catch and hold prey. — **Mantodea** (mantids)




5e. Front wings short and leathery. Hind wings membranous. Elongated flattened body with distinct forceps-like pinchers (cerci). — **Dermaptera** (earwigs)



(4b) 6a. Front wings thickened at base but membranous and overlapping at tips, so that the wings form a triangle pattern on the back. Beak-like mouthparts usually easily visible and appears to arise from front of head, ahead of eyes. — **Hemiptera** (true bugs)




6b. Wings membranous folded tent-like at rest. Beak-like mouthparts not very visible and appear to arise from the area between the front pair of legs. — **Homoptera** (leafhoppers, treehoppers)




(3b) 7a. Wings membranous — go to 8


7b. Wings usually covered with powdery-like scales. Mouthparts in the form of a coiled up tube beneath the head. — *Lepidoptera* (butterflies, moths)




7c. Wings very narrow and fringed with hair. Tiny insect (less than 1/8 inch). Foot ending in balloon-like swelling. — *Thysanoptera* (thrips)



7d. Wings covered with fine hairs, held roof-like over body at rest. Looks like moth with hairy wings. Long antennae extended back over body. — *Trichoptera* (caddisfly)



(7a) 8a. Piercing-sucking mouthparts. Beak-like mouthparts not very visible and appear to arise from the area between the front pair of legs. Wings generally held roof-like over body when at rest. — *Homoptera* (aphids, cicadas, leafhoppers, psyllids, treehoppers, whiteflies)



aphids



cicada




leafhopper



whitefly

8b. Front wing usually larger than hind wing. Wings may be connected. Chewing or chewing sucking mouthparts. Bee, wasp, hornet like. — *Hymenoptera* (bees, wasps, hornets, wings ants)




8c. Large wings in comparison to petite body. Wings usually with many veins and cross veins, held roof-like over body when at rest. Mouthparts chewing. No cerci (tail-like appendages). — *Neuroptera* (lacewing)




8d. Wings longer than body. Body whitish, looks like a fat ant without a slim waist. Straight and beaded antennae. — *Isoptera* (winged termite)



8e. Tiny (less than 1/8 inch). Wings held roof-like over body when at rest. Veins prominent. Hair-like antennae as long or longer than body. Whitish body. — *Psocoptera* (book lice)



8f. Insect dominated by very large eyes. Wings elongated, highly veined. Slender, needle-like body. Chewing mouthparts used to capture and consume prey in flight. — *Odonata* (**dragonflies and damselflies**)



8g. Elongated wings fold flat over body when at rest. Long thread-like antennae. Two tail-like filaments. — *Plecoptera* (**stonefly**)



8h. Front wings large and triangular shaped. Hind wings small and rounded. Held vertical over body. Two-very long tail-like filaments. — *Ephemeroptera* (**Mayfly**)



(1b) 9a. Piercing-sucking mouthparts. — go to 10




9b. Chewing mouthparts. — go to 11




9c. Rasping-sucking mouthparts. Tiny (less than 1/8 inch), slender insects. Balloon-like swelling on end of foot. Often noticed by scarring and deformation of leaves and flowers where it feeds. — *Thysanoptera* (**thrips**)




(9a) 10a. Bug-like with jointed beak. Mouthparts usually easily visible and appear to arise from front of head, in front of eyes. — *Hemiptera* (**true bugs**)




10b. Bug-like with jointed beak. Mouthparts not very visible and appear to arise from the area between the front pair of legs. — *Homoptera* (**aphids, cicadas, leafhoppers, etc.**)




10c. Tiny, hard bodied, flattened sideways, covered with bristles that project backwards. Large hind legs designed for jumping, fast moving. — *Siphonaptera* (**fleas**)




10d. Small, flattened parasites of animals and people. Forelegs with claw designed to grasp hair or feathers. Head narrower than thorax. — *Anoplura* (sucking lice, including head lice and body lice)




(9b) 11a. Larva-like (caterpillar-like, grub-like, maggot-like). — go to 12



11b. Bug-like with tail-like features (cerci, filaments). — go to 15




11c. Ant-like. — go to 16




11d. Dragon-like. — go to 17



11e. Tiny, bug-like. — go to 18



(11a) 12a. Caterpillar-like: soft bodied, three pair of legs on thorax, fleshy leg-like prolegs on abdomen. — go to 13




12b. Grub-like: head capsule, three pair of legs on thorax, no legs on abdomen. — typical of *Coleoptera* (beetle grub) [details page 313-2]




12c. Legless — go to 14


(12a) 13a. Up to five pair of prolegs on abdomen. — *Lepidoptera* (butterfly and moth caterpillar) [details page 313-12]




13b. Six or more pair of prolegs on abdomen. — *Hymenoptera* (sawfly larva)



(12c) 14a. Maggot-like: no legs, no head capsule, mouth-hooks. — typical of *Diptera* (fly maggot)




- 14b. Legless with enlarged head area. — typical of *Coleoptera* (borer larva) [details page 313-2] and some *Hymenoptera* larva




- (11b) 15a. Elongated flattened body with distinct forceps-like pinchers (cerci). — *Dermaptera* (earwigs)




- 15b. Hind legs designed for jumping. Antennae tread-like. one pair of tail-like cerci on most adults. — *Orthoptera* (grasshoppers, crickets, katydids)




- 15c. Flattened body, long thread-like antennae. — *Blattaria* (cockroaches)




- 15d. Front legs designed for grasping and holding prey. Body elongated. Long, thread-like antennae. — *Mantodea* (mantids)




- 15e. Very elongated, stick-like insect. — *Phasmida*, (walkingstick)



- 15f. Two or three long, tail-like cerci. Fast moving. Small 1/4 to 1/2 inch. — *Zygentomaa* (Silverfish, firebrats)



- 15g. Tiny (1-2 mm) soft-bodied insect. “Springtail” often present, used to jump. — *Collembola* (springtail)



Courtesy Colorado State University